

Amino Acids Specifications / Monographs	page	1 / 2
L-Arginine		
Issued Date: Dec, 4, 2015		

L-Arginine¹

C₆H₁₄N₄O₂: 174.20

L-Arginine, when dried, contains not less than 98.5 percent and not more than 101.0 percent of L-Arginine (C₆H₁₄N₄O₂).

Description

White crystals or crystalline powder.

Freely soluble in water and in formic acid, practically insoluble in ethanol (99.5).

Dissolves in dilute hydrochloric acid and it shows hygroscopic.

Identification

Compare the infrared absorption spectrum of the sample with that of the standard by potassium bromide disc method.

Specifications

Item	Limit	Test
Specific rotation [α] _D ²⁰	+26.9 to +27.9°	AJI TEST 1 [Dried sample, C=8, 6mol/L HCl] ²
State of solution (Transmittance)	Clear and colorless Not less than 98.0%	AJI TEST 2 [1.0g in 10mL of H ₂ O, spectrophotometer, 430nm, 10mm cell thickness]
Chloride (Cl)	Not more than 0.020%	AJI TEST 3 [0.5g, A-1, ref: 0.28mL of 0.01mol/L HCl]
Ammonium (NH ₄)	Not more than 0.02%	AJI TEST 4 [D-1]
Sulfate (SO ₄)	Not more than 0.020%	AJI TEST 5 [0.85g, (1), ref: 0.35mL of 0.005mol/L H ₂ SO ₄]
Iron (Fe)	Not more than 10ppm	AJI TEST 6 [1.0g, A-1 ,ref: 1.0mL of Iron Std. (0.01mg/mL)]
Heavy metals (Pb)	Not more than 10ppm	AJI TEST 7 [1.0g, weakly acidic, (1), ref: 1.0mL of Pb Std. (0.01mg/mL)]
Arsenic (As ₂ O ₃)	Not more than 1ppm	AJI TEST 8 [2.0g, (1), ref: 2.0mL of As ₂ O ₃ Std.]
Related substances	1)Conforms ³ 2)Any unspecified impurity Not more than 0.20% Total impurities Not more than 0.50%	AJI TEST 9 [Test sample: 50μg, S-6-a, control; L-Arg 0.25μg] AJI TEST 26 ⁴
Loss on drying	Not more than 0.30%	AJI TEST 11 [1g, at 105°C for 3 hours]
Residue on ignition (Sulfated)	Not more than 0.10%	AJI TEST 13 [1g, at 550°C to 650°C for 3 hours]

Amino Acids Specifications / Monographs		page	2 / 2
L-Arginine			
Issued Date: Dec, 4, 2015			

L-Arginine¹

Specifications (cont'd)

Item	Limit	Test
Assay	98.5 to 101.0%	AJI TEST 14 [Dried sample, 80mg, (1), 3mL of formic acid, 50mL of acetic acid (100), 0.1mol/L HClO ₄ 1mL=8.710mg C ₆ H ₁₄ N ₄ O ₂]
pH	10.5 to 12.0	AJI TEST 33 [1.0g in 10mL of H ₂ O] ⁵

The test for Endotoxin when the material will be used for manufacturing parenteral products is as follows:

Item	Limit	Test
Endotoxin	Less than 6.0EU/g	AJI TEST 34 [C=1, kinetic-turbidimetric technique]

¹ This product, in terms of actual quality, conforms to USP, EP, and JP.

² Temperature coefficient of $[\alpha]_D^{25}$: -0.04°

³ Any secondary spot in the chromatogram obtained from the Test Solution is less intense than the principal spot in the chromatogram obtained from the Standard Solution: the number of those spots is not more than four and not more than 2.0% of the total impurities found.

⁴ Disregard limit: 0.05%

⁵ Care should be taken during measurement against atmospheric carbon dioxide.

End of document