


Nippon Protein Co., Ltd. Product Specification	Established Date	March 1, 2008	Spec. No.	URE-10018H-3	Page	1/2
	Effective Date	December 11, 2014				

L-Cysteine Hydrochloride Monohydrate*1



L-Cysteine Hydrochloride Monohydrate, when calculated on the dried basis, contains not less than 98.5 percent and not more than 101.0 percent of L-Cysteine Hydrochloride (C₃H₇NO₂S·HCl).

Description: White crystals or crystalline powder; Characteristic odor and strong acid taste. Very soluble in water, and soluble in ethanol (99.5). Dissolves in 6mol/L hydrochloric acid TS.

Identification: Compare the infrared absorption spectrum <NP TEST 35> of the sample with that of the standard by potassium bromide disc method.

Specifications:

Item	Limit	Test Method
Specific rotation [α] ^{20D}	+6.1 to +7.8° +6.0 to +7.1°	NP TEST 1 [sample calculated on the dried basis, C=8, 1mol/L HCl]*2 [sample calculated on the dried basis, C=8, 6mol/L HCl]*3
State of solution (Transmittance)	clear and colorless not less than 98.0%	NP TEST 2 [1.0g in 10mL of H ₂ O, spectrophotometer, 430nm, 10mm cell thickness]
Chloride (Cl)	19.89 to 20.29%	NP TEST 3 [350mg, B-2]
Ammonium (NH₄)	not more than 0.02%	NP TEST 4 [(1)]
Sulfate (SO₄)	not more than 0.020%	NP TEST 5 [0.85g, (1), ref: 0.35mL of 0.005mol/L H ₂ SO ₄]
Iron (Fe)	not more than 10ppm	NP TEST 6 [0.75g, B-1, ref: 0.75mL of Iron Std. (0.01mg/mL)]
Heavy metals (Pb)	not more than 10ppm	NP TEST 7 [1.0g, (3), ref: 1.0mL of Pb Std. (0.01mg/mL)]
Arsenic (As₂O₃)	not more than 1ppm	NP TEST 8 [2.0g, (1), ref: 2.0mL of As ₂ O ₃ Std.]
Related substances	conforms*4	NP TEST 9 [test sample: 50μg, B-1-a, Control: L-Cys·HCl·H ₂ O 0.25μg]*5
Loss on drying	8.50 to 12.00%	NP TEST 11 [1g, in vacuum, P ₂ O ₅ , at room temperature for 20 hours]

Nippon Protein Co., Ltd. Product Specification	Established Date	March 1, 2008	Spec. No.	URE-10018H-3	Page	2/2
	Effective Date	December 11, 2014				

Item	Limit	Test Method
Residue on ignition (sulfated)	not more than 0.10%	NP TEST 13 [1g, at 550°C to 650°C for 3 hours]
Assay	98.5 to 101.0%	NP TEST 16 [sample calculated on the dried basis, 250mg, 0.05mol/L I ₂ 1mL = 15.76mg C ₃ H ₇ NO ₂ S·HCl]
pH	1.5 to 2.0	NP TEST 33 [1.0g in 100mL of H ₂ O]

* 1 : USP

* 2 : Temperature coefficient of $[\alpha]_D^{20}$: -0.03°

* 3 : Temperature coefficient of $[\alpha]_D^{20}$: -0.06°

* 4 : Any secondary spot in the chromatogram obtained from the Test Solution is less intense than the principal spot in the chromatogram obtained from the Standard Solution: the number of those spots is not more than four and not more than 2.0% of total impurities is found.

* 5 : Test Solution: Dissolve 100mg of the sample in N-ethylmaleimide solution (1→50) to make 10mL and stand for 30 minutes.

Standard Solution: Dilute 2mL of Test Solution with water to 100mL. Dilute 5mL of this solution with water to 20mL.

Proceed as directed for procedure under NP TEST9 (Thin-layer chromatography).

End of document