Amino Acids Specifications / Monographs		
L-Proline	page	1/2
Issued Date: Dec, 4, 2015		

L-Proline¹

C₅H₉NO₂: 115.13

L-Proline contains not less than 99.0 percent and not more than 101.0 percent of L-Proline (C₅H₉NO₂), calculated on the dried basis.

Description

White crystals or white crystalline powder; slightly sweet taste.

Very soluble in water and in formic acid, and slightly soluble in ethanol (99.5).

It is deliquescent.

Identification

Compare the infrared absorption spectrum of the sample with that of the standard by potassium bromide disc method.

Specifications

Item	Limit	Test
Specific rotation $[\alpha]_D^{20}$	-84.5 to -86.0°	AJI TEST 1
		[Calculated on the dried basis, $C=4$, H_2O] ²
State of solution	Clear and colorless	AJI TEST 2
(Transmittance)	Not less than 98.0%	[1.0g in 10mL of H ₂ O, spectrophotometer, 430nm, 10mm cell
		thickness]
Chloride (Cl)	Not more than 0.020%	AJITEST 3
		[0.5g, A-1, ref: 0.28mL of 0.01mol/L HCl]
Ammonium (NH ₄)	Not more than 0.02%	AJI TEST 4
		[D-1]
Sulfate (SO ₄)	Not more than 0.020%	AJI TEST 5
		[0.85g, (1), ref: 0.35mL of 0.005mol/L H2SO4]
Iron (Fe)	Not more than 10ppm	AJI TEST 6
		[1.0g, A-1, ref: 1.0mL of Iron Std. (0.01mg/mL)]
Heavy metals (Pb)	Not more than 10ppm	AJITEST7
		[1.0g, (1), ref: 1.0mL of Pb Std. (0.01mg/mL)]
Arsenic (As ₂ O ₃)	Not more than 1ppm	AJI TEST 8
		[2.0g, (1), ref: 2.0mL of As ₂ O ₃ Std.]
Related substances	1)Conforms ³	AJITEST 9
		[Test sample: 50µg, B-6-a, control; L-Pro 0.25µg]
	2)Any unspecified impurity	AJI TEST 26 ⁴
	Not more than 0.20%	
	Total impurities	
	Not more than 0.50%	
Loss on drying	Not more than 0.30%	AJI TEST 11
		[1g, at 105°C for 3 hours]
Residue on ignition	Not more than 0.10%	AJI TEST 13
(Sulfated)		[1g, at 550°C to 650°C for 3 hours]

AJINOMOTO CO,. INC.

Amino Acids Specifications / Monographs		
L-Proline	page	2/2
Issued Date: Dec, 4, 2015		

L-Proline

Specifications (cont'd)

Item	Limit	Test
Assay	99.0 to 101.0%	AJI TEST 14
		[Calculated on the dried basis,120mg, (1), 3mL of formic acid,
		50mL of acetic acid (100), 0.1mol/L HClO ₄ 1mL=11.51mg
		$C_5H_9NO_2$
pН	5.9 to 6.9	AJI TEST 33
		[1.0g in 10mL of H ₂ O]

The test for Endotoxin when the material will be used for manufacturing parenteral products is as follows:

Item	Limit	Test
Endotoxin	Less than 6.0EU/g	AJI TEST 34
		[C=1, kinetic-turbidimetric technique]

¹ This product, in terms of actual quality, conforms to USP, EP, and JP.

End of document

² Temperature coefficient of $[\alpha]_D^t$: 0.0°

Any secondary spot in the chromatogram obtained from the Test Solution is less intense than the principal spot in the chromatogram obtained from the Standard Solution: the number of those spots is not more than four and not more than 2.0% of the total impurities found.

Disregard limit: 0.05%